



Endotoksemi Sonrası Kalp Dokusunda Asimetrik dimetilarjinin ve 3-Nitrotirozin Seviyeleri

Asymmetric Dimethylarginine and 3-Nitrotyrosine Levels of the Heart Tissue after Endotoxemia

Asimetrik Dimetilarjinin ve 3-Nitrotirozin / Asymmetric dimethylarginine and 3-Nitrotyrosine

Yasemin Gülcan Kurt¹, Nurten Turkozkan², Halil Yaman¹, Nihal Şahin²,
Hüsamettin Erdamar¹, Gonca Ozan², Filiz Bircan², Emin Özgür Akgül¹, Tuncer Caycı¹, Mehmet Kemal Erbil¹

¹Department of Medical Biochemistry, Gülhane Military Medical Academy,

²Department of Medical Biochemistry, Gazi University Medical Faculty, Ankara, Turkey.

Özet

Amaç

Endotoksemi sonrası kalp disfonksiyonu patojenezinde oksidatif ve nitrozatif stres önemlidir. Asimetrik dimetilarjinin, arjininin yerine geçerek nitrik oksit üretimini yarışmalı olarak inhibe eder. Kardiyovasküler vakaları öngörmede asimetrik dimetilarjinin ve inflamasyon arasındaki bağlantı yakın zamanda gösterilmiştir. Peroksinitrit nitrik oksit ve süperoksit arasındaki etkileşimden oluşan bir reaksiyon ürünüdür.

Gereç ve Yöntemler

Bu çalışmada endotoksemi (4 mg/kg, 6 saatte) sonrasında asimetrik dimetilarjinin ve L-arjinin seviyeleri ve peroksinitrit oluşumunun stabil ürünü olan 3-nitrotirozin arasındaki ilişki araştırılmıştır. Asimetrik dimetilarjinin ve L-arjinin floresan dedektör, 3-nitrotirozin ise elektrokimyasal dedektörlü yüksek basınçlı sıvı kromatografisi ile ölçülmüştür.

Bulgular

Endotoksin verildikten sonra asimetrik dimetilarjinin ve L-arjinin seviyelerinde değişiklik olmadı ama 3-nitrotirozin seviyeleri istatistiksel olarak anlamlı derecede ($p<0.05$) arttı.

Sonuç

4 mg/kg dozunda endotoksin verildikten 6 saat sonra alınan kalp dokularında asimetrik dimetilarjinin ve peroksinitrit üretimleri arasında bir korelasyon bulunmadı.

Anahtar Sözcükler

Asimetrik Dimetilarjinin, Endotoksemi, 3-Nitrotirozin, Kalp.

Abstract

Aim

Oxidative and nitrosative stress are important in the pathogenesis of heart dysfunction after endotoxemia. Asymmetric dimethylarginine competitively inhibits nitric oxide production by displacing L-arginine. A link between asymmetric dimethylarginine and inflammation has been demonstrated recently in the prediction of cardiovascular events. Peroxynitrite is a reaction product formed by the interaction between nitric oxide and superoxide.

Material and Methods

In this study, the relationship between asymmetric dimethylarginine and L-arginine level and 3-nitrotyrosine, a stable product of peroxynitrite formation was investigated after endotoxemia (4 mg /kg at 6h). Measurement of asymmetric dimethylarginine and L-arginine were accomplished by high pressure liquid chromatography with fluorescent detector. 3-nitrotyrosine was quantified by high pressure liquid chromatography with electrochemical detector.

Results

After administration of endotoxin asymmetric dimethylarginine and L-arginine levels did not change but 3-nitrotyrosine levels statistically significantly increased ($p<0.05$).

Conclusion

We could not detect any link between asymmetric dimethylarginine and peroxynitrite production in the heart tissues after given endotoxin using 4 mg / kg dose and at 6 h.

Keywords

Asymmetric dimethylarginine, Endotoxemia, 3-Nitrotyrosine, Heart.

DOI: 10.4328/JCAM.215 Received: 01.03.2010 Accepted: 21.04.2010 Printed: 01.05.2011 J Clin Anal Med 2011;2(2):24-6

Corresponding Author: Yasemin Gülcan Kurt, GATA Tıbbi Biyokimya AD, 06010, Etlik, Ankara, Türkiye.

Phone: +903123043323 Fax: +903123043300 E-mail: ygkurt@gmail.com

Introduction

Overproduction of nitric oxide (NO) has been contributed to tissue injury in inflammatory conditions. One mechanism by which NO can contribute to tissue injury is by its diffusion-controlled reaction with superoxide (O_2^-) to produce peroxynitrite ($ONOO^-$), a potent oxidant thought to be a key mediator of NO mediated tissue injury in a dose-dependent manner in atherosclerosis, congestive heart failure and other disease states involving inflammatory oxidative stress [1-3]. NO is formed from the semi-essential amino acid L-arginine by the action of three isoforms of the nitric oxide synthase (nNOS, cNOS, iNOS). iNOS produces high amount NO after exposure to proinflammatory cytokines [3-5]. $ONOO^-$ reacts with the phenolic ring of tyrosine to form a stable product, 3 nitrotyrosine (3-NT). 3-NT has been widely used as a marker for the detection of $ONOO^-$ formation in vivo [6-8]. Asymmetric dimethylarginine (ADMA) is an L-arginine analog that serves an endogenous inhibitor of both cNOS and iNOS. ADMA is involved in the pathophysiology of vascular disease and has been demonstrated to be not only a marker of endothelial dysfunction, but also a novel cardiovascular risk factor. ADMA has been reported competitively inhibit NO elaboration by displacing L-arginine from NOS [1, 9, 10]. The levels of ADMA and NO are dependent on the prevailing L-arginine concentrations. Clinical and experimental evidences have suggested that elevation of ADMA can cause a relative L-arginine deficiency, even in the presence of normal L-arginine level. In addition to producing NO, NOS can catalyze O_2^- formation in the presence of low L-arginine level [1,11]. The role of ADMA and L-arginine levels in the regulation of $ONOO^-$ production in the heart after endotoxin administration has not yet been well explored. The aim of our study was to examine the effect of endotoxin administration on the heart levels of L-arginine, ADMA and 3-NT and to investigate the relationship between ADMA, L-arginine level and peroxynitrite generation in guinea pigs.

Material and Methods

Animals and study protocol

Following ethics committee approval, all experiments was performed with 20 male Dunkin Hartley guinea pigs weighing 500- 600 g. The animals were randomly divided into two groups. Group I (n=10) served as a control group and was given only saline solution. Group II (n=10) served as an endotoxemia group and was administrated intraperitoneally lipopolysaccharide (LPS) (Escherichia coli LPS serotype O111:B4, from sigma 4mg/kg.).

Measurement of ADMA and L-arginine

Measurement of ADMA and L-arginine level were accomplished by high pressure liquid chromatography (HPLC) using the method described by Momohora, Chen and Azuma et al. [12,13,14]. 300 mg tissue was homogenized in 3 ml phosphate buffer (pH: 6.5). The homogenate was centrifuged at 10000 g for 20 min at 4 OC. Then 60 mg tricarboxylic acid was added to the 1 ml supernatant and the mixture was left in an ice-bath for 10 min. The precipitated protein removed by centrifugation at 3000 g. for 15 min. The supernatant which was filtered through a 0.2 μ m filter was mixed with 100 μ L of derivatization reagent (prepared by dissolving 10 mg o-phthalaldehyde in 0.5 ml of methanol, 2 ml 0.4 M borate buffer (pH:10.0) and 30 μ L of 2-mercaptoethanol) and then injected into the chro-

matographic system. Separation of ADMA and L-arginine levels were achieved with 150 x 4 mm I.D. Nova-pak C18 column with a particle size of 5 μ m (waters, Milipore, Milford MA, USA) using 50 mM sodium acetate (pH:6.8), methanol and tetrahydrofurane as mobile phase (A, 82:17:1; B, 22:77:1) at flow rate 1.0 ml/min. The areas of peaks detected fluorescent detector (Ex: 338 nm; Em: 425) were used for quantification ADMA and L-arginine levels in heart. The analytical variability of the method was less then 7%, and the detection limit of the assay was 0.1 μ M.

Measurement of Tyrosine Nitration

Tissue homogenenates were prepared according to the method described by Kamisaki et al [15]. Briefly, 0.5 g heart tissue was homogenized in 1.5 ml of 50 mM potassium phosphate buffer (pH: 7.2) and then centrifuged for 5 min at 600 g. Following the acid hydrolysis of the precipitate, it was evaporated under nitrogen. After 1 ml distilled water adding, 10 μ L of sample was applied to HPLC. All samples were analysed by HPLC with ECD using the method described by Maruyama et al [16]. For 3-NT detection a Macrotech C-18 analytical column (50 x 1 mm) was used with 50 mM phosphoric acid / 50 mM citric acid / %5 (v/v) methanol (pH:3.1) as the mobile phase. Applied potentials were -850 mV for the reduction and 600 mV for the oxidation detection cell.

Statistical analysis

Results were expressed as mean \pm standard deviation (SD). Statistical analyses were performed using a software program (SPSS 11.5 for Windows, Chicago, IL, USA). The nonparametric Mann-Whitney U test was used to analyse the significance of the difference between control and experimental groups. For tests of significance a p value of less than 0.05 was considered to be significant.

Results

Endotoxin administration did not change levels of ADMA in heart tissues compared to the control group. After endotoxemia levels of L-arginine in heart tissues were reduced but were not statistically significant. 3-NT levels in the heart tissues significantly increased ($p < 0.05$) after endotoxin given. The levels of ADMA, L-Arginine

Table 1. The levels of ADMA, L-Arginine and 3-NT in heart tissues.

	ADMA (μ mol /g tissue)	L-Arginine (nmol /g tissue)	3-NT (nmol /g tissue)
Control group	0.46 \pm 0.0061	37.2 \pm 6.8	0.94 \pm 0.21
Endotoxin group	0.48 \pm 0.094	33.5 \pm 4.3	1.24 \pm 0.17*

Results were expressed as mean \pm standard deviation (SD)
* p <0.05

and 3-NT in heart tissues are shown in Table 1.

Discussion

Recent studies suggested that exposure animal hearts or cardiac myocytes to bacterial endotoxin (LPS) or proinflammatory cytokines enhanced NO level [2, 4, 6, 17]. In cells NO arises from the enzymatic oxidation (by NO synthase) of guanidino nitrogen of one terminal of L-arginine. $ONOO^-$ is reaction product formed by the interaction between NO and O_2^- and it has also been implicated as a toxic mediator in the pathophysiology of endotoxemia in atherosclerosis [1, 2]. $ONOO^-$ reacts with tyrosine to form, a stable product, 3-NT [3, 6]. ADMA, a naturally occurring analog of L-arginine, competitively inhibits NO production by NOS. L-arginine is an important substrate for both NO and ADMA production. A link between ADMA and inflammation has been demonstrated in the

prediction of cardiovascular events [18]. We examined the relationship between ONOO- production and levels of ADMA, L-arginine in heart tissues after endotoxin administration. The present study for the first time demonstrated increased 3-NT level in endotoxemic heart tissues, but concurrent with unchanged ADMA production. In addition to this study provides evidence for decreased L-arginine level but no significant. In a previously our study we found elevated plasma ADMA levels in same animals with endotoxemia, as well as decreased L-arginine levels and declined the ratio of Larginine/ADMA, but NO levels did not change [19]. In contrast to the plasma, in this study, levels of ADMA and L-arginine in heart tissues did not change. But after endotoxemia in the heart tissues of same animals, ONOO- production was high. It may be dependent on the mechanism found in the heart exposed to LPS. Our results may be consistent with previous observation. Investigators have demonstrated that the 3-NT levels were increased in LPS-treated hearts (4 mg/kg). The other investigators examined the cytotoxic effect of

iNOS-generated NO in cultured cardiac myocytes treated with LPS. They showed that the cytokines caused nitrotyrosine formation in the injured myocytes [2-4]. In our study, while the levels of 3-NT in animal heart tissues increased with endotoxemia, ADMA and L-arginine levels decreased but this decrease was not significant. There was no correlation between ADMA, L-arginine and 3-NT. These results were in agreement with previous observations in vitro that ADMA preferentially inhibited cNOS isoform, while having minimal effect on NO production by iNOS [1]. In the other in vivo study, it has been reported that E. coli mediated endotoxemia decreases the plasma arginine level but ADMA did not change [19]. Ueda et al. demonstrated exogenous administration of NO donors dose-dependently increased NO synthesis in the cultured media but had no effect on ADMA and L-arginine levels [20]. In conclusion we could not find a link between ADMA, L-arginine concentrations and ONOO- production in the heart after endotoxin administration. More researches may be needed to elucidate these results using different dose and time after LPS administration.

References

1. Wells SM, Holian A. Asymmetric dimethylarginine induces oxidative and nitrosative stress in murine lung epithelial cells. *Am J Respir Cell*. 2007;36:520-8.
2. Usui, M.; Matsuoka, H.; Miyazaki, H.; Ueda, S.; Okuda, S.; Imaizumi, T. Increased endogenous nitric oxide synthase inhibitor in patients with congestive heart failure. *Life Sci*. 1998;62:2425-30.
3. Keira N, Tatsumi T, Matoba S, Shiraishi J, Yamanaka S, Akashi K. et al. Lethal effect of cytokine induced nitric oxide and peroxynitrite on cultured rat cardiac myocytes. *J Mol Cell Cardiol*. 2002;34(5):583-96.
4. Khadour FH, Panas D, Ferdinandy P, Schulze C, Csont T, Lalu MM, et al. Enhanced NO and superoxide generation in dysfunctional hearts from endotoxemic rats. *Am J Physiol Heart Circ Physiol*. 2002;283(3):H1108-15.
5. Yin GQ, Du KH, Gu FR, Fang ZX, Tang JQ, Zhong B, et al. Early-phase endotoxemic shock-induced myocardial injury increases iNOS and selectin expression in macaque primate. *Heart Lung Circ*. 2007;16(2):85-92.
6. Iqbal M, Cohen RI, Marzouk K, Liu SF. Time course of nitric oxide, peroxynitrite, and antioxidants in the endotoxemic heart. *Crit Care Med*. 2002;30(6):1291-6.
7. Cimen B, Turkozkan N, Unlu A, Erbil MK. Effects of melatonin on 3-nitrotyrosine formation and energy charge ratio in guinea pig kidney in LPS-induced stress. *Cell Biochem Funct*. 2005;23(4):273-7.
8. Seven I, Turkozkan N, Cimen B. The effects of nitric oxide synthesis on the Na⁺/K⁺-ATPase activity in guinea pig kidney exposed to lipopolysaccharides. *Mol Cell Biochem*. 2005;271(1-2):107-12.
9. Chan JR, Böger RH, Bode-Böger SM, Tangphao O, Tsao P S, Blaschke TF, et al. Asymmetric dimethylarginine increase mononuclear adhesiveness hypercholesterolemic humans. *Arterioscler Thromb Vasc Biol*. 2000;20(4):1040-6.
10. Nijveldt RJ, Siroen MP, Teerlink T, van Lambalgen AA, Rauwerda JA, van Leeuwen PA. Gut and liver handling of asymmetric and symmetric dimethylarginine in the rat under basal conditions and during endotoxemia. *Liver Int*. 2004;24 (5): 510-8.
11. Jia YX, Pan CS, Yang JH, Liu XH, Yuan WJ, Zhao J, et al. Altered L-arginine/nitric oxide synthase/nitric oxide pathway in the vascular adventitia of rats with sepsis. *Clin Exp Pharmacol Physiol*. 2006;33(12):1202-8.
12. Momohara Y, Sakamoto S, Obayashi S, Aso T, Goto M, Azuma H. Roles of endogenous nitric oxide synthase inhibitors and endothelin-1 for regulating myometrial contractions during gestation in the rat. *Mol Hum Reprod*. 2004;10(7):505-12.
13. Chen BM, Xia LW, Zhao RQ. Determination of N(G),N(G)-dimethylarginine in human plasma by high-performance liquid chromatography. *J Chromatogr B Biomed Sci Appl*. 1997;692(2): 467-71.
14. Azuma H, Masuda H, Sato J, Niwa K, Tokoro T. A possible role of endogenous inhibitor for nitric oxide synthesis in the bovine ciliary muscle. *Exp Eye Res*. 1997;64(5):823-30.
15. Kamisaki Y, Wada K, Nakamoto K, Kishimoto Y, Kitano M, Itoh T. Sensitive determination of nitrotyrosine in human plasma by isocratic high-performance liquid chromatography. *J Chromatogr B Biomed Appl*. 1996;685(2):343-7.
16. Maruyama W, Hashizume Y, Matsubara K, Naoi M. Identification of 3-nitro-L-tyrosine, a product of nitric oxide and superoxide, as an indicator of oxidative stress in the human brain. *J Chromatogr B Biomed App*. 1996;676(1):153-8.
17. Matsushita A, Iwase M, Kato Y, Ichihara S, Ichihara G, Kimata H, et al. Differential cardiovascular effects of endotoxin derived from *Escherichia coli* or *Pseudomonas aeruginosa*. *Exp Anim*. 2007;56(5):339-48.
18. Zoccali C, Benedetto FA, Maas R, Mallamaci F, Tripepi G, Malatino LS, et al. Asymmetric dimethylarginine, C-reactive protein, and carotid intima-media thickness in end-stage renal disease. *J Am Soc Nephrol*. 2002;13(2):490-6.
19. Balabanli B, Erdamar H, Turkozkan N, Yaman H, Kurt Y. Effect of taurine on endotoxin-induced alterations in plasma asymmetric dimethylarginine, L-arginine and nitric oxide in guinea pigs. *J Thromb Thrombolysis*. 2007;24(1):53-7.
20. Ueda S, Kato S, Matsuoka H, Kimoto M, Okuda S, Morimatsu M, et al. Regulation of cytokine-induced nitric oxide synthesis by asymmetric dimethylarginine: role of dimethylarginine dimethylaminohydrolase. *Circ Res*. 2003;92(2):226-33.